arXiv:1207.5770v1 [astro-ph.SR] 24 Jul 2012

EXPLAINING THE TYPE Ia SUPERNOVA PTF 11kx WITH THE CORE DEGENERATE SCENARIO

Noam Soker[1], Amit Kashi[2], Enrique García–Berro[3,4], Santiago Torres[3,4], Judit Camacho[3,4]

ABSTRACT

We argue that the multiple shells of circumstellar material (CSM) and the supernovae (SN) ejecta interaction with the CSM starting 59 days after the explosion of the Type Ia SN (SN Ia) PTF 11kx, are best described by the core-degenerate (CD) scenario for SN Ia. In the CD scenario the super-Chandrasekhar mass white dwarf (WD) is formed at the termination of the common envelope phase from a merger of a WD companion with the hot core of a massive asymptotic giant branch (AGB) star. In most cases the WD is destructed and accreted onto the more massive core. However, in rare cases where mergers take place when the WD is denser than the core, the core will be destructed and accreted onto the cooler WD. In such cases the explosion might occur with no appreciable delay, i.e., months to years after the termination of the common envelope (CE) phase. This, we propose, is the evolutionary route that lead to the explosion of PTF 11kx. The CD scenario can account for the very massive CSM within ~ 1000 AU of the exploding PTF 11kx star, for the presence of hydrogen, and for the presence of shells in the CSM.

1. INTRODUCTION

Observations and theoretical studies cannot teach us yet whether all three scenarios for the formation of Type Ia supernova (SN Ia), or only one or two of them can work — e.g., Livio (2001), Maoz (2010), Howell (2011). These three basic theoretical scenarios can be

---

[1] Department of Physics, Technion – Israel Institute of Technology, Haifa 32000, Israel; soker@physics.technion.ac.il

[2] Department of Physics & Astronomy, University of Nevada, Las Vegas, 4505 S. Maryland Pkwy, Las Vegas, NV, 89154-4002, USA

[3] Departament de Física Aplicada, Universitat Politècnica de Catalunya, c/Esteve Terrades 5, 08860 Castelldefels, Spain

[4] Institute for Space Studies of Catalonia, c/Gran Capità 2–4, Edif. Nexus 104, 08034 Barcelona, Spain

described as follows. (*i*) In the single-degenerate (SD) scenario (e.g., Whelan & Iben 1973; Nomoto 1982; Han & Podsiadlowski 2004) a WD grows in mass through accretion from a non-degenerate stellar companion. (*ii*) In the double-degenerate (DD) scenario (Webbink 1984; Iben & Tutukov 1984; see van Kerkwijk et al. 2010 for a paper on sub-Chandrasekhar mass remnants) two WDs merge after losing energy and angular momentum through the radiation of gravitational waves. (*iii*) In the core-degenerate (CD) scenario for the formation of SN Ia the Chandrasekhar or super-Chandrasekhar mass WD is formed at the termination of the CE phase, or during the planetary nebula phase, from a merger of a WD companion with the hot core of a massive AGB star (Kashi & Soker 2011; Ilkov & Soker 2012a; Soker 2011). The merger of a WD with the core of an AGB star was studied in the past (Sparks & Stecher 1974; Livio & Riess 2003; Tout et al. 2008). Livio & Riess (2003) suggested that the merger of the WD with the AGB core leads to a SN Ia that occurs at the end of the CE phase or shortly after, and can explain the presence of hydrogen lines. Due to its rapid rotation (e.g., Anand 1965; Ostriker & Bodenheimer 1968; Uenishi et al. 2003; Yoon & Langer 2005; Lorén-Aguilar et al. 2009), and possibly very strong central magnetic fields (e.g., Kundu & Mukhopadhyay 2012), the super-Chandrasekhar WD explosion might be substantially delayed.

The recently observed SN Ia PTF 11kx (Dilday et al. 2012) has narrow lines, including hydrogen lines, and indications of interaction with the a massive circumstellar medium (CSM) which starts 59 days after the explosion. Dilday et al. (2012) argued that PTF 11kx can be explained by the SD-scenario for SN Ia. They dismiss the CD scenario on several grounds, like that it cannot account for several CSM shells. We find this unjustified in section 2.

The destruction of the WD and its accretion onto the core while the core is still hot might prevent an early ignition of carbon (Yoon et al. 2007), which is one of the theoretical problems of the DD scenario (e.g., Saio & Nomoto 2004). However, in the case of PTF 11kx ignition of the merger product is required quite early, within ~ 30 yr from the CE ejection. We therefore consider in section 3 the possibility that in the case of PTF 11kx the core was destructed onto the cooler WD. In cases when the hot core is destructed onto the cooler WD and explosion occurs shortly after merger. Thus, there is no delay by spin-down or emission of gravitational waves, and the CD and the DD scenario overlap. In section 4 we show that these cases can eject a massive envelope and account for the frequency of such SN Ia. Our summary is in section 5.

2. THE CIRCUMSTELLAR MEDIUM (CSM)

Starting 59 days after the explosion, Dilday et al. (2012) detected a group of absorption lines (Na I, Fe II, Ti II, and He I) and Hα and Hβ P-cygni lines at a velocity shift of 65 km s^{-1}. Ca II H&K absorption line was detected with outflow velocity of 100 km s^{-1}. A group of emission lines (H, He, Fe, Ti) at 100 km s^{-1} as well, and Dilday et al. (2012) suggested they arrive from the same location as the Ca II H&K absorption line. This, according to Dilday et al. (2012), indicates the presence of multiple CSM shells interacting with the SN ejecta. Adopting an ejecta velocity of 25 000 km s^{-1} they deduced that the ejecta-CSM interaction starts at a distance of $\sim 10^{16}$ cm.

Dilday et al. (2012) suggest that PTF 11kx is a type Ia SN formed by the SD scenario. A claim that was raised also in the theoretical work of Hachisu et al. (2012). In the scenario of Dilday et al. (2012) the CSM structure is assumed to originate from recent recurrent nova eruptions whose ejecta sweeps the wind of a giant companion. Based on the saturated Ca II HK absorption lines Dilday et al. (2012) calculated the CSM mass to be $M_{\rm CSM} \simeq 5.3kM_\odot$, where k is the covering fraction. Since the SD scenario cannot account for such a high mass, as the wind from a giant star cannot explain such a mass within $\sim$ 1000 AU, they argue that $k \ll 1$ and the mass is much lower than $5M_\odot$, but give no specific number. However, to obtain a saturated line the absorbing gas should cover the exploding sphere, and hence it is required that $k > 0.25$. We conclude, based on the analysis of Dilday et al. (2012), that $M_{\rm CSM} > 1M_\odot$.

SN 2002ic (Hamuy et al. 2003) was also classified as a SN Ia, but with strong and broad, FWHM of > 1000 km s^{-1}, Hα emission. The Hα emission suggested that it arose from an interaction between the SN ejecta and a dense H-rich CSM of a mass of $\sim 6M_\odot$ (Wang et al. 2004). Nomoto et al. (2005) — see also Nomoto et al. (2004) — developed a model to account for the light curve and suggested that the CSM mass is $\sim 1.3M_\odot$ and is non spherical or very clumpy. Livio & Riess (2003) proposed that SN 2002ic represents a rare case in which the explosion occurs during (or immediately following) the CE phase, and the critical mass limit is reached by core-WD merger. Livio & Riess (2003) considered this scenario to be a subtype of the DD scenario. However, a merger while the core is still hot and there is no time delay due to gravitational waves emitted by a double-degenerate remnant is now termed the CD scenario (Kashi & Soker 2011; Ilkov & Soker 2012a), as it has some fundamental differences from the DD scenario (Soker 2011).

SN 2005gj is another member of SN Ia with hydrogen emission (Aldering et al. 2006, Prieto et al. 2007), and an aspherical or clumpy CSM (Aldering et al. 2006). As days from the explosion passed, more emission lines of the hydrogen Balmer series and He I lines were detected gradually, what may indicate the presence of a few layers of CSM. Aldering et al.

(2006) estimated the mass in an Hα shell to be $\sim 0.016 - 0.16 M_\odot$, but the total mass might be larger.

The SNe above, and specifically PTF 11kx, can be explained in the frame of the CD scenario, as we expect three main mass loss episodes: (1) Before the CE. This is the ordinary mass loss of the AGB but enhanced by tidal interaction. (2) The CE ejection. At this stage the spiral structure can form several shells along the line of sight. Some numerical simulations — e.g., Ricker & Taam (2012); Sandquist et al. (1998) — show that the mass loss from a CE process occurs in more than one episode, i.e., there are a few peaks in the mass loss rate. In particular, along each direction there can be several shells due to the spiral structure that is formed during the CE phase, and that can later have imprint in the CSM. These shell would contain all together the mass of the AGB envelope when the CE started, so each shell would have $\sim 0.5 - 3 M_\odot$. (3) During the merger. This mass loss event would eject $\sim 0.1 - 0.5 M_\odot$. It can accelerate some previously ejected gas to higher velocities, e.g., ~ 100 km s^{-1}.

3. THE CASE FOR A MASSIVE CORE-WD MERGER

When two WD merge to form a super-Chandrasekhar mass WD a huge amount of energy is released. Actually, this might give an event brighter than expected from SN Ia explosion, which is one of the problems of the DD scenario. In the present case, the merger is of a core of an AGB star and a WD companion (the remnant of the primary star), with masses of $M_{\rm core}$ and $M_{\rm WD}$, respectively. The final state just before the SN Ia explosion is that the primary accreted a mass ΔM from the core, and the rest of the mass is expelled. The difference in binding energy is between the energy required to unbind the core and the energy released by the accretion of mass onto the WD:

$$\Delta E_G \simeq 0.5 \frac{G(M_{\rm WD} + 0.5\Delta M)\Delta M}{R_{\rm WD}} - \Gamma \frac{G M_{\rm core}^2}{R_{\rm core}}, \tag{1}$$

where $\Gamma \simeq 0.5$ is a parameter that depends on the properties of the core. We assume an adiabatic index of $\gamma = 5/3$ as the core is hot, and took a coefficient of 0.5 in the first term of equation (1). For the typical parameters used here the gravitational energy released is

$$\begin{aligned} \Delta E_G({\rm erg}) \simeq 1.9 \times 10^{50} \left(\frac{M_{\rm WD} + 0.5\Delta M}{1.3 M_\odot}\right) \left(\frac{\Delta M}{0.6 M_\odot}\right) \left(\frac{R_{\rm WD}}{5400 \text{ km}}\right)^{-1} \\ - 1.3 \times 10^{50} \left(\frac{\Gamma}{0.5}\right) \left(\frac{M_{\rm core}}{1 M_\odot}\right)^2 \left(\frac{R_{\rm core}}{10^4 \text{ km}}\right)^{-1}. \end{aligned} \tag{2}$$

The radius of a WD of mass $1 M_\odot$ is 6000 km (Provencal et al. 1998). But as matter is accreted onto the cold massive WD its radius shrinks. However, this will not release much energy as the matter in the center is relativistic, and the adiabatic index γ is close to the full relativistic value 4/3. For $\gamma = 4/3$, the marginally-stable value, no energy is released as the star contracts in hydrostatic equilibrium. Here it is not fully relativistic, but the adiabatic index is still below 5/3 and not much energy will be released. The binding energy of a WD at the critical mass of $1.4 M_{\odot}$ is $E_b = -5.5 \times 10^{50}$ erg (e.g., Ibanez Cabanell 1984). For a merger WD mass of $1.4 M_{\odot}$ and $\gamma = 5/3$ the corresponding radius is 4700 km. We conclude that not much energy will be released by the contraction of the accreting cold WD, and took an effective radius of 5400 km.

The conclusion is that the total energy that has to be radiated *before* the explosion is $E_{\rm pre} \simeq$ several $\times 10^{49}$ erg. This is very sensitive to the parameters due to the subtraction of two very close values. For example, adopting $M_{\rm WD} = 1.1 M_{\odot}$ and $\Delta M = 0.45 M_{\odot}$, and using the same formalism we find the released energy to be $\sim 10^{49}$ erg. The total energy released in radiation can be larger than that emitted during the SN explosion itself! However, the model assumes (something that needs verification) that the building of the merger remnant to the point it explodes lasts for several years. The time scale is two orders of magnitudes longer than that of the SN explosion. The bolometric luminosity will be over one order of magnitude lower than the SN explosion, and most of the radiation (due to the optically thick ejected envelope) be emitted in the IR. This is the reason it avoided detection by different surveys in the visible. However, the model predicts that such SN Ia might have precursor lasting several years with 5–10 bolometric magnitudes fainter than the SN explosion, and peaks in the near IR. In case the merger is of a lighter core-WD system, then the pre-explosion total energy released might be much larger. But in that case the delay will be longer, as the central region needs to cool to reach the critical limit for explosion.

4. THE EVOLUTIONARY ROUTE

For the progenitor of the explosion we look for systems having the following properties.

1. The total mass of the secondary core during the final CE phase and the mass of the WD remnant of the primary star should be super-Chandrasekhar, $M_{\rm core} + M_{\rm WD} > 1.4 M_{\odot}$. But as discussed in the previous section, to prevent a pre-SN explosion outburst, the core and WD should be much more massive, $M_{\rm core} + M_{\rm WD} \gtrsim 2 M_{\odot}$.

2. The total mass of the envelope at the final CE phase must be as massive as the observed CSM. We take it to be $M_{\rm env} \gtrsim 2 M_{\odot}$, but prefer $M_{\rm env} \gtrsim 4 M_{\odot}$.

3. To facilitate merger at the termination of the CE phase we require $M_{\rm env}/M_{\rm WD} \gtrsim 3$, but prefer a larger number even (Soker 2012).

4. It is likely that an explosion without a delay will take place when the core is accreted onto the cooler WD remnant of the primary star (Kashi & Soker 2011). For that the density of the core should be lower than the density of the WD. As the core is hot its radius is larger than that of a cool WD. The condition reads $M_{\rm core} \lesssim 1.5 M_{\rm WD}$.

5. As we require the merger product to be a CO WD, to avoid ONe WDs, we are limited to both $M_{\rm core} < 1.1 M_\odot$ and $M_{\rm WD} < 1.1 M_\odot$.

To examine for such systems we run the population synthesis code described in García-Berro et al. (2012), where all details are given. Here, for the sake of conciseness, we only present the results for systems that obey the constraints above. In Figure 1 we plot the envelope mass of the secondary at the time of merger in the plane $M_{\rm core} - M_{\rm WD}$, while in Figure 2 we present the ratio of the envelope mass to the WD mass. Note that the likely progenitors of PTF 11kx are within a region of WD masses between ~ 0.9 and $\sim 1.1 M_\odot$, while the masses of the hydrogen-exhausted core are also within ~ 0.9 and $\sim 1.1 M_\odot$. The most restrictive conditions are, respectively, conditions 1 and 3. Namely, we find that the progenitors of this supernova must fulfill that the total mass of the merged remnant should be larger than $2 M_\odot$, and also that the ratio of the envelope mass to the mass of the white dwarf should be larger than ~ 3, although other possibilities cannot be presently discarded.

Dilday et al. (2012) crudely estimate that the fraction of SNe Ia that exhibit prominent circumstellar interaction near maximum light is $\sim 0.1 - 1\%$. As $\sim 1 - 2$ SN Ia occur per $1000 M_\odot$ stars formed (Maoz et al. 2012), the estimate of Dilday et al. (2012) stands at $\sim 0.001 - 0.02$ SN Ia with massive CSM per $1000 M_\odot$ stars formed. Our Monte Carlo simulations show that the frequency of these systems with the conditions listed above is 0.002 per $1000 M_\odot$ stars formed. If we relax the assumption on the core+WD mass and the envelope mass to $M_{\rm core} + M_{\rm WD} \gtrsim 1.8 M_\odot$ (instead of $M_{\rm core} + M_{\rm WD} \gtrsim 2 M_\odot$), and $M_{\rm env} \gtrsim 0.5 M_\odot$ (instead of $M_{\rm env} \gtrsim 4 M_\odot$), respectively, we find 0.016 systems to obey the condition per $1000 M_\odot$ stars formed. The later envelope mass might be appropriate to SN 2005gj. We conclude that the CD scenario can satisfactorily explain all the observed properties of PTF 11kx and, moreover, that the number of expected progenitors with massive CSM is consistent with the observations.

5. SUMMARY

The conclusion that the presence of any CSM around a SN Ia implies its association with the single-degenerate (SD) scenario — e.g., Sternberg et al. (2011) — is problematic, in particular in cases where the CSM mass within ~ 1000 AU is $\gtrsim 0.01 M_\odot$. An explosion set by the merger of a WD companion with the core of the giant star naturally occurs within a massive CSM — the ejected common envelope (Livio & Riess 2003). The formation of super Chandrasekhar WDs through the merger of a WD with the core at the termination of the CE phase as an explanation for most (or even all) SN Ia is termed the core-degenerate scenario (CD). The CD scenario can occur for a large range of WD masses. For the majority of the systems the WD is destructed and accreted onto the more massive core. For a specific, narrow range of WD and envelope masses, a SN Ia explosion with H-rich environment like PTF 11kx can occur.

We find the association of the SN Ia PTF 11kx (Dilday et al. 2012) with the SD scenario to be unlikely due to the massive CSM (Dilday et al. 2012). Instead, we found that the merger of a massive WD with a massive core, as marked by the thick line on the upper right corner of Figures 1 and 2, to be able to account for the properties of PTF 11kx. We also found from our population synthesis calculations that the number of systems with large total mass of the WD and core match the number of SN Ia with massive CSM as deduced from observations. Adding the recent finding of Ilkov & Soker (2012b) from a population synthesis study that the CD scenario can account for most, or even all, SNe Ia, we conclude that the CD scenario is a main channel for the formation of SNe Ia.

This research was supported by the Asher Fund for Space Research at the Technion, by MCINN grant AYA2011–23102, by the European Union FEDER funds, and by the ESF EUROGENESIS project (grant EUI2009-04167).

REFERENCES

Aldering, G., Antilogus, P., Bailey, S., et al. 2006, ApJ, 650, 510

Anand, S. P. S. 1965, Proceedings of the National Academy of Science, 54, 23

Dilday, B., Howell, D. A., Cenko, S. B., et al. 2012, arXiv:1207.1306

García-Berro, E., Lorén-Aguilar, P., Aznar-Siguán, G., et al. 2012, ApJ, 749, 25

Hachisu, I., Kato, M., & Nomoto, K. 2012, arXiv:1207.2261

Hamuy, M., Phillips, M. M., Suntzeff, N. B., et al. 2003, Nature, 424, 651

Han, Z., & Podsiadlowski, P. 2004, MNRAS, 350, 1301

Howell, D. A. 2011, Nature, Communications, accepted, arXiv:1011.0441

Ibanez Cabanell, J. M. 1984, A&A, 135, 382

Iben, I., Jr., & Tutukov, A. V. 1984, ApJS, 54, 335

Ilkov, M., & Soker, N. 2012a, MNRAS, 419, 1695

Ilkov, M., & Soker, N. 2012b, submitted

Kashi, A., & Soker, N. 2011, MNRAS, 417, 1466

Kundu, A., & Mukhopadhyay, B. 2012, arXiv:1204.1463, Modern Physics Letters A, in press

Livio, M. 2001, Supernovae and Gamma-Ray Bursts: the Greatest Explosions since the Big Bang, eds. Mario Livio, Nino Panagia, Kailash Sahu. Space Telescope Science Institute symposium series, Vol. 13. Cambridge University Press, (Cambridge, UK) 334

Livio, M., & Riess, A. G. 2003, ApJ, 594, L93

Lorén-Aguilar, P., Isern, J., & García-Berro, E. 2009, A&A, 500, 1193

Maoz, D. 2010, American Institute of Physics Conference Series, 1314, 223

Maoz, D., Mannucci, F., & Brandt, T. D. 2012, arXiv:1206.0465

Nomoto, K. 1982, ApJ, 253, 798

Nomoto, K., Suzuki, T., Deng, J., et al. 2004, Frontier in Astroparticle Physics and Cosmology, 323

Nomoto, K., Suzuki, T., Deng, J., Uenishi, T., & Hachisu, I. 2005, 1604-2004: Supernovae as Cosmological Lighthouses, 342, 105

Ostriker, J. P., & Bodenheimer, P. 1968, ApJ, 151, 1089

Prieto, J. L., Garnavich, P. M., Phillips, M. M., et al. 2007, arXiv:0706.4088

Provencal, J. L., Shipman, H. L., Hog, E., & Thejll, P. 1998, ApJ, 494, 759

Ricker, P. M., & Taam, R. E. 2012, ApJ, 746, 74

Saio, H., & Nomoto, K. 2004, ApJ, 615, 444

Sandquist, E. L., Taam, R. E., Chen, X., Bodenheimer, P., & Burkert, A. 1998, ApJ, 500, 909

Soker, N. 2011, arXiv:1109.4652

Soker, N. 2012, arXiv:1204.3173

Sparks, W. M., & Stecher, T. P. 1974, ApJ, 188, 149

Sternberg, A., Gal-Yam, A., Simon, J. D., et al. 2011, Science, 333, 856

Tout, C. A., Wickramasinghe, D. T., Liebert, J., Ferrario, L., & Pringle, J. E. 2008, MNRAS, 387, 897

Uenishi, T., Nomoto, K., & Hachisu, I. 2003, ApJ, 595, 1094

van Kerkwijk, M. H., Chang, P., & Justham, S. 2010, ApJ, 722, L157

Wang, L., Baade, D., Höflich, P., et al. 2004, ApJ, 604, L53

Webbink, R. F. 1984, ApJ, 277, 355

Whelan, J., & Iben, I., Jr. 1973, ApJ, 186, 1007

Yoon, S.-C., & Langer, N. 2005, A&A, 435, 967

Yoon, S.-C., Podsiadlowski, P., & Rosswog, S. 2007, MNRAS, 380, 933

This preprint was prepared with the AAS LaTeX macros v5.2.



Fig. 1.— Mass of the envelope of the giant as a function of the mass of the primary white dwarf and of the mass of the hydrogen-exhausted core of the secondary. The different color scales denote different masses of the envelope. The brown scale corresponds to relatively small masses of the envelope ($2 \leq M_{\rm env}/M_\odot \leq 3$), the gray scale corresponds to intermediate masses ($3 \leq M_{\rm env}/M_\odot \leq 4$), while the blue scale has been selected to show the masses of interest for the scenario described in the main text ($4 \leq M_{\rm env}/M_\odot \leq 5$). Also plotted are the constrains given by conditions 1 and 4. For the sake of clarity, the region corresponding to the likely progenitors of PTF 11kx has been highlighted.



Fig. 2.— Ratio of the mass of the envelope of the secondary to the mass of the primary white dwarf, as given by the colored bar, as a function of the mass of the primary white dwarf and of the hydrogen-exhausted core of the secondary. The rest of the lines are the same shown in Figure 1. Again, the region of likely progenitors of PTF 11kx has been conveniently highlighted in this plane.